

# Circular ensembles

September 29, 2006

Dyson introduced the circular ensemble of eigenvalues of a Haar distributed unitary matrix that has pdf  $\prod_{j < k} |e^{i\theta_j} - e^{i\theta_k}|^2$ . A generalization of this is the pdf

$$\prod_{j < k} |e^{i\theta_j} - e^{i\theta_k}|^2 \cdot \prod_{k=1}^n e^{-V(\theta_k)}.$$

These are closely related to orthogonal polynomials w.r.t. the weight  $e^{-V(\theta)}$  on the unit circle. Basic facts can be found in Mehta's book (1).

Riemann-Hilbert techniques apply naturally to the study of scaling limits of the spacings in the case when  $V$  is analytic. How to go beyond analytic  $V$ ?

Miller and McLaughlin (2) have a generalization of R-H techniques for non-analytic  $V$  and apply it to asymptotics of orthogonal polynomials w.r.t  $e^{-V}$ . These techniques have not been applied to the study of correlation functions yet. Specific projects:

1. Concrete question: How many derivatives of  $V$  are needed to deduce sine kernel correlations?
2. If  $V$  is  $k$  times continuously differentiable but the  $(k + 1)^{\text{st}}$  derivative has a jump discontinuity at a point, the scaled correlations at that point may have a different kernel which should form a universality class of its own? This would be a new phenomenon. See also the website <http://www.math.lsa.umich.edu/millerpd/ResearchAndPublications/CP.html>

## References

- [1] Mehta, Madan Lal. (2004). Random matrices. Third edition. Pure and Applied Mathematics (Amsterdam), 142. Elsevier/Academic Press, Amsterdam, 2004.
- [2] McLaughlin, K. T.-R. and Miller, P. D. (2006). The  $\bar{\partial}$  steepest descent method and the asymptotic behavior of polynomials orthogonal on the unit circle with fixed and exponentially varying nonanalytic weights. *Int. Math. Res. Pap.* Art. ID 48673, 1–77. arxiv:math.CA/0406484